

Citation for published version:

Pipoly, I, Bokony, V, Kirkpatrick, M, Donald, PF, Székely, T & Liker, A 2015, 'The genetic sex-determination system predicts adult sex ratios in tetrapods', *Nature*, vol. 527, no. 7576, pp. 91-94.
<https://doi.org/10.1038/nature15380>

DOI:

[10.1038/nature15380](https://doi.org/10.1038/nature15380)

Publication date:

2015

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**The genetic sex-determination system predicts
adult sex ratios in tetrapods**

Ivett Pipoly¹, Veronika Bókony^{1,2}, Mark Kirkpatrick³, Paul F. Donald⁴, Tamás Székely^{5*} &
András Liker^{1*}

¹ Department of Limnology, University of Pannonia, Pf. 158, H-8201 Veszprém, Hungary

² Lendület Evolutionary Ecology Research Group, Plant Protection Institute, Centre for
Agricultural Research, Hungarian Academy of Sciences, Herman Ottó út 15, H-1022
Budapest, Hungary

³ Department of Integrative Biology, University of Texas, 78712, Austin, Texas, USA

⁴ RSPB Centre for Conservation Science, RSPB, The Lodge, Sandy, Bedfordshire SG19 2DL,
UK

⁵ Department of Biology and Biochemistry, Biodiversity Laboratory, University of Bath, BA2
7AY Bath, United Kingdom

*These authors contributed equally to this work.

Adult sex ratio (ASR) has critical effects on behaviour, ecology and population dynamics¹⁻³, but the causes of variation in ASRs are unclear^{4,5}. Here we assess whether the type of genetic sex determination influences ASR using data from 344 species in 117 families of tetrapods. We find that taxa with female heterogamety have a significantly more male-biased ASR (mean proportion of males: 0.55 ± 0.01 SE) than taxa with male heterogamety (0.43 ± 0.01). The genetic sex-determination system explains 24% of interspecific variation in ASR in amphibians and 36% in reptiles. We consider several genetic factors that could contribute to this pattern, including meiotic drive and sex-linked deleterious mutations, but further work is needed to quantify their effects. Regardless of the mechanism, the effects of the genetic sex-determination system on the adult sex ratio are likely to have profound impacts on the demography and social behaviour of tetrapods.

Adult sex ratio (ASR) varies widely in nature, ranging from populations that are heavily male-biased to ones composed only of adult females⁵⁻⁸. Birds and schistosome parasites tend to have male-biased ASR, for example, while mammals and copepods usually exhibit female-biased ASR⁵. Extreme bias occurs among marsupials (Didelphidae and Dasyuridae): males die after the mating season, so there are times when the entire population consists of pregnant females⁹. Understanding the causes and consequences of ASR variation is an important goal in evolutionary biology, population demography and biodiversity conservation because ASR impacts behaviour, breeding systems, and ultimately population fitness^{1,2,10-13}. It is also a significant issue in social sciences, human health and economics, since unbalanced ASRs have been linked to violence, rape, mate choice decisions and spread of diseases like HIV¹⁴⁻¹⁶. The causes of ASR variation in wild populations, however, remain obscure^{5,13,17}.

One factor that could impact the ASR is the genetic sex-determination system^{7,8,18}. Taxa such as mammals and fruit flies have XY sex determination (males are heterogametic), whereas taxa such as birds and butterflies have ZW sex determination (females are heterogametic). Sex-determination systems could affect the ASR in several ways. A skewed ASR might result from an unbalanced sex ratio at birth caused by sex ratio distorters¹⁹. Alternatively, a biased ASR could develop after birth if sex chromosomes contribute to sex differences in mortality^{8,18,20,21}. Differential postnatal mortality is likely to be the main driver of biased ASR in birds and mammals, since birth sex ratios in these classes tend to be balanced⁷.

Here we use data from the four major clades of tetrapods (amphibians, reptiles, birds, and mammals) to assess whether adult sex ratios differ between taxa with XY and ZW sex determination (Fig. 1; Supplementary Table 1). While mammals and birds are fixed for XY and ZW sex determination, respectively, reptiles and amphibians provide particularly attractive opportunities for this study since transitions between sex-determination systems have occurred many times within these clades^{22,23}. We compiled published data on adult sex ratios in wild populations and their sex-determination systems (Supplementary Table 1). To control for phylogenetic effects, we used phylogenetic generalized least squares (PGLS)²⁴ to test for differences in ASRs between XY and ZW taxa, and Pagel's discrete method (PDM)²⁵ to test whether XY and ZW systems are evolutionarily associated with female-biased and male-biased sex ratios, respectively. Phylogenies were taken from recent molecular studies (see Methods for details).

Both the ASR and the sex-determination system are highly variable across tetrapods (Fig. 1, Supplementary Table 1). We find that ASR and sex determination are correlated. Before controlling for phylogenetic effects, we find that ASRs are significantly more male-biased in species with ZW sex determination than in those with XY sex determination (Fig. 2, Table 1, Extended Data Table 1). Similarly, the proportion of species with male-biased ASRs is greater among ZW than among XY species (Fig. 1, Table 1). These differences are significant within amphibians, within reptiles, and across tetrapods as a whole (Table 1, Extended Data Table 1).

The pattern remains significant after controlling for phylogenetic effects. Both the mean of ASR across species (analyzed using PGLS) and the proportion of species with male-biased sex ratios (analyzed using PDM) are significantly different between XY and ZW systems within amphibians, within reptiles, and across tetrapods as a whole (Table 1, Extended Data Table 1). The effect is strong in clades with variation in sex determination: the type of genetic sex determination explains up to 24% of the interspecific variance in ASR among amphibians and 36% in reptiles (estimated using PGLS, Extended Data Table 2). The results remain significant when we treat three large clades with invariant sex-determination systems as a single datum each (snakes, ZW; birds, ZW; mammals, XY; Extended Data Table 1), when we make different assumptions about branch lengths in the phylogeny (Extended Data Table 2), and when we use arc-sine transformed ASR values and control for variance in sample size (see Methods).

Body size and breeding latitude correlate with life-history traits in many organisms and these traits could affect ASR^{26–28}. Sexual size dimorphism is linked to differential sexual selection acting on males and females and thus influences sex-specific mortality, and has

been suggested to drive the evolution of genetic sex-determination systems²⁹.
Nevertheless, we find that neither body size nor breeding latitude explains the ASR in
phylogenetically controlled multi-predictor analyses (Table 2). Sexual size dimorphism is
significantly associated with ASR in reptiles and across tetrapods as a whole, but the
effect of the genetic sex-determination system remains significant when size dimorphism
is included in the analysis (Table 2).

Sex differences in dispersal may also result in biased ASRs. However, dispersal is
unlikely to explain the relationship between ASR and sex-determination systems. First,
male-biased dispersal is typical in reptiles regardless of sex-determination system
(Supplementary Material 1)^{30,31}. Second, there is no relationship between ASR and sex
bias in dispersal distance in birds (Supplementary Material 1). Finally, the relationship
between sex determination and ASR remain significant when the influence of sex-biased
dispersal is controlled for in multi-predictor models in tetrapods (Supplementary Material
1).

The sex-determination system may affect the ASR in the directions seen in the data in a
number of ways. First, sexual selection can fix mutations that increase male mating
success and decrease male survival. They will accumulate on Y but not W chromosomes,
and will accumulate more readily on X than Z chromosomes if they tend to be recessive.
Second, biased ASRs could result from recessive mutations at loci carried on the X (or Z)
but absent from the Y (or W) chromosome since they are not masked in the heterogametic
sex (the “unguarded sex chromosome” hypothesis)^{7,8,18}, and from deleterious mutations
carried on the Y (or W) but not on the X (or Z). At loci carried on both sex chromosomes,
alleles on the Y (or W) can show partial degeneration³². Population genetic models

suggest deleterious mutation pressure alone may not be adequate to explain ASR biases as large as those observed (Supplementary Material 2), but the models do not include factors that could be important, notably degeneration of Y and W chromosomes by genetic drift³². A third hypothesis is imperfect dosage compensation, which may be deleterious to the heterogametic sex³³. Fourth, distorted sex ratios can result from meiotic drive acting on sex chromosomes^{34,35}. Drive more often produces female-biased sex ratios in XY systems at birth³⁶. There is little data on drive in ZW systems, but if it operates in a symmetric fashion then we expect it to cause male-biased sex ratios. Fifth, Y and W chromosomes might degenerate during the lifespan, for example by telomere shortening or loss of epigenetic marks, more rapidly than X and Z chromosomes. A final possibility is that sex-antagonistic selection acting on sex-linked loci could lead to biased sex ratios, but unlike the preceding hypotheses there does not seem to be a robust prediction about the direction of the ASR bias it will produce (see Supplementary Material 2).

The limited data that are available do not provide clear support for any of these hypotheses, although critical tests are lacking. For instance, the meiotic drive process predicts biased sex ratios at birth. Although a recent comparative analysis in birds suggests that sex ratios at birth are unrelated to biased ASRs¹⁰, offspring sex ratios have not been compared between different sex-determination systems. Additional insight might come from study of dioecious plants with biased sex ratios³⁷, but their skewed ASR could result from selection on the gametophytic stage that is absent from animals³⁸.

Evolutionary feedbacks from the ASR to the sex-determination system are also possible: for example, the ASR could influence sexual size dimorphism and sexual conflict, which in turn could trigger transitions in sex determination^{29,39,40}.

In conclusion, we demonstrate strong and phylogenetically robust associations between genetic sex-determination systems and a demographic property of populations, ASR. Although the mechanisms that drive this association need further theoretical and empirical analyses, the observed pattern is biologically important for two reasons. First, changes in sex-determination systems are expected to have knock-on effects on social behaviour. Theory suggests that ASR affects violence, pair bonds, infidelity and parental care^{1,41}, and field-based studies support these predictions^{4,13,15,16}. For instance, female-biased ASRs co-occur with polygyny and female care, whereas male-biased ASRs tend to co-occur with polyandry and male care in birds⁴. Second, sex-determination systems may have important demographic consequences through skewed birth sex ratios and sex-biased survival. Such biases may not only impact upon the productivity and growth of populations, but also their genetic composition and viability. Further theoretical, experimental, and comparative studies are clearly needed to understand the linkages between sex determination, demography, and social behaviour.

Methods Summary

We collected ASR data for 39 amphibian, 67 reptile, 187 bird and 51 mammal species from the literature. When more than one estimate was available for a species we used their mean. Because genetic sex-determination systems of amphibians vary between closely related species⁴², we used only those species in which sex determination was characterized at the species level^{22,43}. Sex determination is evolutionarily less labile in reptiles, thus we included all species for which sex determination was known either at the family level or, in variable families, at the species level^{22,44,45}. Breeding latitude was calculated as the distance from the Equator in latitudinal degrees (averaged if multiple records were

available for a species). Sexual dimorphism in body size was calculated as $\log_{10}(\text{male size}) - \log_{10}(\text{female size})$. All data and their sources are given in Supplementary Table 1.

We controlled for phylogenetic effects in two ways. First, we tested whether ASR bias (female- or male-biased) is evolutionary associated with the type of sex determination using Pagel's discrete method (PDM)²⁵ as implemented in *BayesTrait*. Second, we tested for differences in ASR with phylogenetic generalized least squares (PGLS)²⁴ using the R package *caper*⁴⁶. We used recently published phylogenies for amphibians⁴⁷, reptiles^{48–50}, birds⁵¹, and mammals^{52,53}. The branching topology between these four major clades was based on recent tetrapod phylogenies^{54,55} (Fig. 1).

We developed simple population genetic models of the effects that deleterious mutation and sex-antagonistic selection can have on the ASR (Supplementary Material 2). The results regarding deleterious mutations reported in the text assume the mutations are largely or entirely recessive and have multiplicative fitness effects across loci, that the loci are in linkage equilibrium, and that selection is strong relative to mutation and drift. The loci are assumed to be fully sex-linked, and those carried on the one type of sex chromosome (e.g. the X) are assumed to have no homologue on the other type (e.g. the Y) that could otherwise mask a deleterious mutation. Fitness effects of mutations in hemizygotes and homozygotes are assumed equal. In our models of sex-antagonistic selection the loci are fully sex-linked and carried on an X or Z chromosome. The models for both deleterious mutation and sex-antagonistic selection are deterministic. They therefore do not account for stochastic processes (such as “Muller’s ratchet”) that are thought important to sex chromosome evolution³². It is possible that the conclusions would change qualitatively if the models were extended to include those additional effects.

References

1. Kokko, H. & Jennions, M. D. Parental investment, sexual selection and sex ratios. *J. Evol. Biol.* **21**, 919–948 (2008).
2. Le Galliard, J.-F., Fitze, P. S., Ferrière, R. & Clobert, J. Sex ratio bias, male aggression, and population collapse in lizards. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 18231–18236 (2005).
3. Fitze, P. S. & Le Galliard, J. F. Operational sex ratio, sexual conflict and the intensity of sexual selection. *Ecol. Lett.* **11**, 432–439 (2008).
4. Liker, A., Freckleton, R. P. & Székely, T. The evolution of sex roles in birds is related to adult sex ratio. *Nat. Commun.* **4**, 1587 (2013).
5. Székely, T., Weissing, F. J. & Komdeur, J. Adult sex ratio variation: implications for breeding system evolution. *J. Evol. Biol.* 1–13 (2014).
6. Breitwisch, R. Mortality patterns, sex ratios, and parental investment in monogamous birds. *Curr. Ornithol.* **6**, 1–50 (1989).
7. Donald, P. F. Adult sex ratios in wild bird populations. *Ibis* **149**, 671–692 (2007).
8. Trivers, R. L. Parental investment and sexual selection. in *Sex. Sel. Descent Men* (B, C.) (Aldine, 1972).
9. Cockburn, A., Scott, M. P. & Dickman, C. R. Sex ratio and intrasexual kin competition in mammals. *Oecologia* **66**, 427–429 (1986).
10. Székely, T., Liker, A., Freckleton, R. P., Fichtel, C. & Kappeler, P. M. Sex-biased survival predicts adult sex ratio variation in wild birds. *Proc. R. Soc. B* **281**, (2014).
11. Bessa-Gomes, C., Legendre, S. & Clobert, J. Allee effects, mating systems and the extinction risk in populations with two sexes. *Ecol. Lett.* **7**, 802–812 (2004).
12. Veran, S. & Beissinger, S. R. Demographic origins of skewed operational and adult sex ratios: perturbation analyses of two-sex models. *Ecol. Lett.* **12**, 129–143 (2009).
13. Liker, A., Freckleton, R. P. & Székely, T. Divorce and infidelity are associated with skewed adult sex ratios in birds. *Curr. Biol.* **24**, 880–884 (2014).
14. Griskevicius, V. *et al.* The financial consequences of too many men: sex ratio effects on saving, borrowing, and spending. *J. Pers. Soc. Psychol.* **102**, 69–80 (2012).
15. Schacht, R., Rauch, K. L. & Borgerhoff Mulder, M. Too many men: the violence problem? *Trends Ecol. Evol.* **29**, 214–222 (2014).

- 222 16. Kandrik, M., Jones, B. C. & DeBruine, L. M. Scarcity of female mates predicts
223 regional variation in men's and women's sociosexual orientation across US states. *Evol.*
224 *Hum. Behav.* **36**(3), 206-210 (2015).
- 225 17. Wilson, E. O. *Sociobiology: The New Synthesis*. (Harvard University Press, 1975).
- 226 18. Haldane, J. B. Sex-ratio and unisexual sterility in hybrid animals. *J. Genet.* **12**, 101–
227 109 (1922).
- 228 19. Burt, A. & Trivers, R. *Genes in Conflict - The Biology of Selfish Genetic Elements*.
229 Harvard University Press, 632 (2008).
- 230 20. Promislow, D. E. L., Montgomerie, R. & Martin, T. E. Mortality costs of sexual
231 dimorphism in birds. *Proc. R. Soc. B* **250**, 143–150 (1992).
- 232 21. Liker, A. & Székely, T. Mortality costs of sexual selection and parental care in natural
233 populations of birds. *Evolution*. **59**, 890–897 (2005).
- 234 22. Ashman, T.-L. *et al.* Tree of Sex Consortium: Tree of Sex: A database of sexual
235 systems. *Sci. data* (2014). at <<http://treeofsex.org/>>
- 236 23. Bachtrog, D. *et al.* Tree of Sex Consortium: Sex determination: why so many ways of
237 doing it? *PLoS Biol.* **12**, (2014).
- 238 24. Pagel, M. Inferring evolutionary processes from phylogenies. *Zool. Scr.* **26**, 331–348
239 (1998).
- 240 25. Pagel, M. Detecting correlated evolution on phylogenies: A general method for the
241 comparative analysis of discrete characters. *Proc. R. Soc. B* **255**, 37–45 (1994).
- 242 26. Du, W., Robbins, T. R., Warner, D. A., Langkilde, T. & Shine, R. Latitudinal and
243 seasonal variation in reproductive effort of the eastern fence lizard (*Sceloporus*
244 *undulatus*). *Integr. Zool.* **9**, 360–371 (2014).
- 245 27. Iverson, J. B., Balgooyen, C. P., Byrd, K. K. & Lyddan, K. K. Latitudinal variation in
246 egg and clutch size in turtles. *Can. J. Zool.* **71**, 2448–2461 (1993).
- 247 28. Morrison, C. & Hero, J.-M. Geographic variation in life-history characteristics of
248 amphibians: a review. *J. Anim. Ecol.* **72**, 270–279 (2003).
- 249 29. Adkins-Regan, E. & Reeve, H. K. Sexual dimorphism in body size and the origin of
250 sex-determination systems. *Am. Nat.* **183**, 519–536 (2014).
- 251 30. Myers, E. A. *et al.* Multilocus phylogeographic assessment of the California Mountain
252 Kingsnake (*Lampropeltis zonata*) suggests alternative patterns of diversification for the
253 California Floristic Province. *Molecular Ecology*. **22**, 5418–5429 (2013).

- 254 31. Qi, Y., Yang, W., Lu, B. & Fu, J. Genetic evidence for male-biased dispersal in the
255 Qinghai toad-headed agamid *Phrynocephalus vlangalii* and its potential link to
256 individual social interactions. *Ecology and Evolution* **3**(5): 1219–1230 (2013).
- 257 32. Bachtrog, D. A dynamic view of sex chromosome evolution. *Curr. Opin. Genet. Dev.*
258 **16**, 578–585 (2006).
- 259 33. Mank, J. E. Sex chromosome dosage compensation: definitely not for everyone. *Trends*
260 *Genet.* **29**, 677–683 (2013).
- 261 34. Hurst, L. D. & Pomiankowski, A. Maintaining mendelism: Might prevention be better
262 than cure? *BioEssays* **13**, 489–490 (1991).
- 263 35. Jaenike, J. Sex chromosome meiotic drive. *Annu. Rev. Ecol. Syst.* **32**, 25–49 (2001).
- 264 36. Werren, J. H. & Beukeboom, L. W. Sex determination, sex ratios, and genetic conflict.
265 *Annu. Rev. Ecol. Evol. Syst.* **29**, 233–261 (1998).
- 266 37. Field, D. L., Pickup, M. & Barrett, S. C. H. Comparative analyses of sex-ratio variation
267 in dioecious flowering plants. *Evolution*, **67**(3), 661–672 (2012).
- 268 38. Hough, J., Immler, S., Barrett, S. C. H. & Otto, S. P. Evolutionarily stable sex ratios
269 and mutation load. *Evolution*, **67**(7), 1915–1925 (2013).
- 270 39. Roberts, R. B., Ser, J. R. & Kocher, T. D. Sexual conflict resolved by invasion of a
271 novel sex determiner in Lake Malawi Cichlid fishes. *Science*. **326**, 998–1001 (2009).
- 272 40. van Doorn, G. S. & Kirkpatrick, M. Transitions between male and female heterogamety
273 caused by sex-antagonistic selection. *Genetics* **186**, 629–645 (2010).
- 274 41. McNamara, J. M., Székely, T., Webb, J. N. & Houston, A. I. A dynamic game-theoretic
275 model of parental care. *Journal of Theoretical Biology* **205**, 605–623 (2000).
- 276 42. Miura, I., Ohtani, H. & Ogata, M. Independent degeneration of W and Y sex
277 chromosomes in frog *Rana rugosa*. *Chromosom. Res.* **20**, 47–55 (2012).
- 278 43. Evans, B. J., Pyron, R. A. & Wiens, J. J. Polyploidization and Sex Chromosome
279 Evolution in Amphibians, in *Polyploidy Genome Evol.* (Soltis, P. S. & Soltis, D. E.)
280 Springer Berlin Heidelberg, 385–410 (2012).
- 281 44. Sarre, S. D., Ezaz, T. & Georges, A. Transitions between sex-determining systems in
282 reptiles and amphibians. *Annu. Rev. Genomics Hum. Genet.* **12**, 391–406 (2011).
- 283 45. Pokorná, M. & Kratochvíl, L. Phylogeny of sex-determining mechanisms in squamate
284 reptiles: are sex chromosomes an evolutionary trap? *Zool. J. Linn. Soc.* **156**, 168–183
285 (2009).
- 286 46. Orme, A. D. *et al.* 'caper': Comparative Analyses of Phylogenetics and Evolution in R.
287 R package version 0.5. (2013).

47. Pyron, R. A. & Wiens, J. J. A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of extant frogs, salamanders, and caecilians. *Mol. Phylogenet. Evol.* **61**, 543–583 (2011).
48. Gardner, M. G., Hugall, A. F., Donnellan, S. C., Hutchinson, M. N. & Foster, R. Molecular systematics of social skinks : phylogeny and taxonomy of the Egernia group (Reptilia : Scincidae). *Zool. J. Linn. Soc.* **154**, 781–794 (2008).
49. Pyron, R. A., Burbrink, F. T. & Wiens, J. J. A phylogeny and revised classification of Squamata, including 4161 species of lizards and snakes. *BMC Evol. Biol.* **13**, 93 (2013).
50. Guillon, J.-M., Guery, L., Hulin, V. & Girondot, M. A large phylogeny of turtles (Testudines) using molecular data. *Contrib. to Zool.* **81**, (2012).
51. Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K. & Mooers, A. O. The global diversity of birds in space and time. *Nature* **491**, 444–448 (2012).
52. Meredith, R. *et al.* Impacts of the cretaceous terrestrial revolution and KPg extinction on mammal diversification. *Science* **334**, 521–524 (2011).
53. Fritz, S. A., Bininda-Emonds, O. R. P. & Purvis, A. Geographical variation in predictors of mammalian extinction risk: big is bad, but only in the tropics. *Ecol. Lett.* **12**, 538–549 (2009).
54. Chiari, Y., Cahais, V., Galtier, N. & Delsuc, F. Phylogenomic analyses support the position of turtles as the sister group of birds and crocodiles (Archosauria). *BMC Biol.* **10**, 65 (2012).
55. Amemiya, C. T. *et al.* The African coelacanth genome provides insights into tetrapod evolution. *Nature* **496**, 311–316 (2013).

Supplementary Information is linked to the online version of the paper at

www.nature.com/nature.

Acknowledgements

Matt Pennell and György Imreh helped constructing the phylogeny figure. We thank D. Bachtrog, K. Reinhold and three reviewers for helpful suggestions. We were supported by the European Union, with the co-funding of the European Social Fund (I.P. by TÁMOP-4.2.2.A-11/1/KONV-2012-0064), and by the U.S. National Science Foundation (DEB-0819901 to

M.K.). TS was supported by a Humboldt Award and MTA-DE “Lendület” grant in projects that lead to the current work. A.L. was supported by the Hungarian Scientific Research Fund (OTKA K112838) and a Marie Curie Intra-European Fellowship.

Author Contributions

T.S., A.L. and V.B. designed the study. I.P., V.B., P.F.D. and A.L. collected the reptile, amphibian, mammal and bird data, respectively. I.P., V.B. and A.L. conducted the analyses. M.K. developed the population genetic models. All authors wrote the paper.

Author Information

Reprints and permissions information are available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of this article at www.nature.com/nature. Comments and requests for materials should be addressed to A.L. (aliker@almos.uni-pannon.hu).

Table 1. The effect of the sex-determination system on the adult sex ratio.

Taxon	Number of species	Mean ASR				% species with male-biased ASR		
		XY	ZW	<i>t</i> -test [†]	PGLS [†]	XY	ZW	PDM [†]
Amphibians	39	0.51	0.61	**	**	42.9	90.9	*
Reptiles	67	0.45	0.57	***	***	24.2	76.5	*
Birds	187	--	0.55	--	--	--	76.5	--
Mammals	51	0.37	--	--	--	9.8	--	--
Tetrapods	344	0.43	0.55	***	***	22.3	77.2	***

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, -- no data or not tested

[†]Detailed results of the statistical analyses are presented in Extended Data Table 1.

Mean adult sex ratios (ASR, proportion of males in the population), *t*-tests and percentage of species with male-biased ASRs represent species-level statistics and analyses, while phylogenetic generalized least squares (PGLS)²⁴ and Pagel's discrete method (PDM)²⁵ were used for phylogenetically corrected analyses of the difference in ASR between XY and ZW species.

Table 2. The relationships between adult sex ratio, sex-determination system and other factors in phylogenetically corrected multi-predictor analyses.

	Amphibians (<i>n</i> = 39)			Reptiles (<i>n</i> = 67)			Tetrapods (<i>n</i> = 259)		
	<i>b</i> (± SE)	<i>t</i>	<i>P</i>	<i>b</i> (± SE)	<i>t</i>	<i>P</i>	<i>b</i> (± SE)	<i>t</i>	<i>P</i>
Sex-determination system	0.10 (± 0.03)	3.38	0.002	0.10 (± 0.02)	4.56	<0.001	0.10 (± 0.02)	5.23	<0.001
Body size	0. (± 0)	1.41	0.166	0 (± 0)	0.78	0.440	0 (± 0)	0.05	0.962
Breeding latitude	0 (± 0)	0.13	0.898	0 (± 0)	0.04	0.966	0 (± 0)	0.24	0.811
Sexual size dimorphism	-0.32 (± 0.34)	0.92	0.363	-0.31 (± 0.15)	2.17	0.034	-0.38 (± 0.07)	5.57	<0.001

Results of phylogenetic generalized least squares (PGLS)²⁴. Separate models were constructed for amphibians, reptiles and tetrapods, that included ASR as response variable. For sex determination, *b* is the estimated difference in ASR between ZW and XY species.

Figure 1. Phylogenetic distribution of adult sex ratio (ASR) and genetic sex-determination systems across tetrapods. Inner band shows the type of sex determination (red: XY, blue: ZW) and the outer band shows the ASR bias for each species included in the study (red: ≤ 0.5 , blue: > 0.5). Sample sizes: 39 species for amphibians, 67 species for reptiles, 187 species for birds and 51 species for mammals (see Supplementary Table 1).

Figure 2. Variation in adult sex ratio as a function of the sex-determination system in amphibians, reptiles, mammals and birds, and tetrapods (all four clades combined). Central dots and solid whiskers are means ± 1 SE, horizontal bars are medians, and boxes and dashed whiskers show the interquartile ranges and data ranges, respectively, based on species values. Numbers of species are at the bottom of each panel. See Table 1 and Extended Data Table 1 for statistical results and Extended Data Figure 1 for phylogenetically corrected graphs.

Methods

Data collection

We collected data on ASR (proportion of males in the adult population) in amphibians and reptiles from literature published by December 2013, by searching in Google Scholar and Web of Science with the key words "sex ratio" and "reptile" or "amphibian" or the scientific names of species. We also used reviews to identify additional data sources^{43,56}. ASR data for mammals⁷ were obtained from a similar search finished in 2007; and we used avian ASR estimates from our existing data set (Supplementary Information of Liker et al.¹³).

We specifically collected ASR data for amphibians and reptiles from studies that aimed to obtain representative estimates for the population composition and thus provide reliable sex ratio data⁵⁷. These include either long-term demographic studies applying mark-recapture or sacrificing methods (i.e. each individual was counted only once) with similar capture probabilities for the sexes, or total population counts. When more than one measure was available, we used the total counts of individually marked animals over the study period because this may best approximate the overall ASR. We excluded studies in which the authors explicitly stated or speculated that their data may not represent the population-level ASR, or when the methods were not described in enough detail to assess the reliability of the ASR estimate. Moreover, we tested whether ASR estimates differ between sampling (hand-capture, trap, other) and marking (mark-recapture, sacrifice) methods, and we found no such differences (linear mixed-effects model with species as random factor, sampling: $F_{(3, 105)} = 0.50$, $P = 0.683$; marking: $F_{(2, 105)} = 2.18$, $P = 0.118$; $n = 234$ records). When more than one estimate of ASR was available for the same population (e.g. from several yearly counts at the same location) we took their mean weighted by sample size. When more than one independent record was available for a species from different populations or studies, we used

their simple mean. Weighted and non-weighted mean ASRs were highly correlated (amphibians: Pearson's $r = 0.973$, $P < 0.001$, $n = 35$ species; reptiles: $r = 0.995$, $P < 0.001$, $n = 60$ species); we used non-weighted averages because not all studies reported sample size.

We categorized the genetic sex-determination (GSD) systems of the species from published sources either as male-heterogametic (XY) or female-heterogametic (ZW). For amphibians, only species with known GSD system were included^{22,43}, because GSD is an evolutionarily labile trait in amphibians; species within a genus or even populations within a species can differ in GSD⁴². For reptiles, we included all species for which GSD was known either at the family level, or at the species level if both XY and ZW systems were reported in the family^{22,44,45}. Our result for reptiles is not changed qualitatively by restricting our analyses to those species for which GSD is known at species level²², i.e. when species for which we assumed GSD based on other species in the family were excluded (difference between XY and ZW reptile species, phylogenetic generalized least squares model (PGLS)^{24,58}: $b \pm SE = 0.11 \pm 0.02$; $t = 4.70$, $P < 0.001$, $n = 26$; $R^2 = 0.479$). All birds were assigned to ZW, and all mammals to XY sex-determination systems²².

We also collected data on three additional ecological and behavioural variables to control for their known correlation with ASR and so reduce potential confounding effects in multi-predictor analyses. First, we used body size (in mm) which was measured as snout to vent length for amphibians and squamates, and carapace length for the two turtle species, where possible from the same population for which ASR was reported. Head-body length was used for mammals ($n = 36$)⁵⁹. Since head-body length is not available for the vast majority of birds, we calculated this from the total body length by extracting bill and tail length ($n = 133$;

Supplementary Table 1). Where we had sex-specific data, the mean of male and female head-body length was used as body size variable in the analyses.

Second, we estimated sexual size dimorphism (SSD) as $\log_{10}(\text{male body size}) - \log_{10}(\text{female body size})$. For birds, we used body mass dimorphism (data available for $n = 181$ species)⁶⁰ due to the lack of sex-specific body length data. The results of the multivariate PGLS model of tetrapods presented in Table 2 remain qualitatively the same when wing length dimorphism (data available for $n = 153$ species) is used for birds instead of body mass dimorphism (effect of sex determination: $b \pm SE = -0.10 \pm 0.02$, $t = 4.97$, $P < 0.001$; body size: $b \pm SE = 0 \pm 0$, $t = 0.06$, $P = 0.949$; latitude: $b \pm SE = 0 \pm 0$, $t = 0.223$, $P = 0.823$; size dimorphism: $b \pm SE = -0.52 \pm 0.12$, $t = 4.33$, $P < 0.001$; $n = 248$ species).

Third, we included breeding latitude as the geographic coordinates of the ASR studies for amphibians and reptiles, taking absolute values to represent distance from the Equator in latitudinal degree. When the authors did not report latitude, we used Google Earth to estimate it on the basis of the description of the study site. For birds and mammals, we used the latitudinal midpoint of the breeding range of the species ($n = 182$ and 44 species, for birds and mammals, respectively; sources: V. Remes, A. Liker, R. Freckleton and T. Székely unpublished data for birds, and the PanTHERIA database for mammals⁶¹, respectively). Mean values of these variables were used if multiple data of body size, latitude or size dimorphism per species were available.

Other possible confounding factors include the lifespan of individuals and sex-specific dispersal distances. First, longer average lifespan may lead to exaggeration of ASR bias. However, in species with available data⁶², lifespan is unrelated to ASR (PGLS, birds: $b \pm SE$

= 0 ± 0 , $t = 0.196$, $P = 0.845$, $n = 71$ species; mammals: $b \pm SE = 0 \pm 0$, $t = 0.751$, $P = 0.457$, $n = 35$ species) and also to the absolute deviation of ASR from 0.5 (i.e. when assuming that longer lifespan can exaggerate ASR bias in either direction; birds: $b \pm SE = 0 \pm 0$, $t = 1.543$, $P = 0.127$, $n = 71$ species; mammals: $b \pm SE = 0 \pm 0$, $t = 0.180$, $P = 0.858$, $n = 35$ species). Second, sex-specific dispersal can bias ASR due to the higher mortality in the sex with longer dispersal distances. However, we found no evidence of a relationship of sex bias in dispersal either with GSD in reptiles, or with ASR in birds (Supplementary Material 1). For these reasons, as well as because data on lifespan and/or sex-specific dispersal are not available for most species in our ASR data set, we did not include these variables in the main multi-predictor models (see Supplementary Material 1 for additional models including dispersal).

Our final dataset comprises data on 39 amphibian species and 67 reptile species (in total $n = 229$ ASR records from different populations), 187 bird species and 51 mammalian species (a total of 344 species). We could not find body size and latitude data for some species, thus sample sizes were reduced in multi-predictor models. All species-level data and their sources are given in Supplementary Table 1.

Data analysis

To assess the reliability of the amphibian and reptile ASR estimates, we calculated the repeatability of ASR as the intraclass correlation coefficient (ICC) following Lessells & Boag⁶³, using only those species for which we had at least two ASR estimates from different populations. These analyses show a moderate repeatability of ASR, and that a significant part of ASR variation is interspecific (amphibians: $ICC = 0.559$, $F_{(22,96)} = 7.27$, $P < 0.001$, $n = 23$ species, $n = 120$ records; reptiles: $ICC = 0.524$, $F_{(13,26)} = 4.11$, $P = 0.001$, $n = 14$ species, $n = 40$ records). For birds, our earlier analyses showed that nearly half (44%) of the ASR

variation was interspecific, and that the direction of ASR (i.e. male- or female-biased) was highly conserved: in 44 species out 55 (80%), the direction of ASR bias was the same for all repeated estimates⁵. For mammals, we did not find enough multiple ASR data within species to estimate repeatability.

In the comparative analyses we used the topology of Pyron & Wiens⁴⁷ for amphibians, a composite phylogeny for reptiles^{48–50}, Jetz et al.⁵¹ for birds¹³, the family-level relationships of Meredith et al.⁵² and the genus/species level relationships of Fritz et al.⁵³ for mammals. For analyses across tetrapods, the branching topology between these four major clades was based on recent tetrapod phylogenies^{54,55} (Fig. 1). Since we did not have branch length information for these composite phylogenies, we ran the analyses using arbitrary gradual branch lengths according to Nee's method⁶⁴. However, our results remained consistent when we repeated the analyses with other branch length assumptions (Pagel's method and unit branch lengths⁶⁴; Extended Data Table 2).

To test the association between ASR bias (male- versus female-biased) and GSD (XY versus ZW) in phylogenetically corrected analyses, we used Pagel's discrete method²⁵ as implemented in *BayesTrait*⁶⁵. We used maximum likelihood methods to fit independent and dependent models for transitions in ASR bias and GSD states, and compared the fit of these two models by a likelihood ratio test²⁵. To test the ASR difference between XY and ZW species, we used PGLS models with maximum likelihood estimates of Pagel's lambda values²⁴ using the R⁶⁶ package *caper*^{46, 58}. ASR was the response variable in all models, and genetic sex-determination system was fitted as the predictor (Table 1, Extended Data Table 1). The parameter estimate b shows the difference in ASR (proportion of males in the population) between ZW and XY species. To test the robustness of the bivariate results, we

490 added body size, breeding latitude and SSD as predictors in multi-predictor models to control
491 for their potential confounding effects (Table 2). As in earlier ASR studies^{5,7}, the distribution
492 of ASR values did not deviate significantly from normal in the four clades separately as well
493 as in tetrapods as a whole; our results remain qualitatively identical when ASR is arc-sine
494 transformed before PGLS analyses (amphibians: $b \pm SE = 0.10 \pm 0.03$, $t_{37} = 3.44$, $P = 0.001$, n
495 $= 39$; reptiles: $b \pm SE = 0.12 \pm 0.02$, $t_{65} = 5.95$, $P < 0.001$, $n = 67$; tetrapods: $b \pm SE = 0.11 \pm$
496 0.02 , $t_{342} = 5.24$, $P < 0.001$, $n = 344$).

497

498 The difference between XY and ZW systems for tetrapods is not sensitive to the inclusion of
499 large clades with uniform sex-determination systems (snakes and birds are all ZW, mammals
500 are all XY) since it remains unchanged when each of these clades is reduced to a single datum
501 of its mean ASR (PGLS: $b \pm SE = 0.10 \pm 0.02$, $t = 5.07$, $P < 0.001$, $R^2 = 0.232$, $n = 87$).

502 Furthermore, our result is also robust to between-species differences in sample size: when we
503 added $\log(\text{no. individuals})$ to the previous model, the effect of sex determination remained
504 significant ($b \pm SE = 0.15 \pm 0.07$, $t = 2.08$, $P = 0.041$) while sample size had no significant
505 effect on ASR ($b \pm SE = 0 \pm 0.01$, $t = 0.35$, $P = 0.72$, $n = 78$). Furthermore, sample size was
506 not a significant predictor of ASR when we added it as a fourth confounding variable in the
507 full PGLS model ($b \pm SE = 0 \pm 0.01$, $t = 1.16$, $P = 0.250$, $n = 78$), and the effect of other
508 predictors remained qualitatively the same as in Table 2. Finally, the results do not change
509 when we only used the most reliable ASR data (based on mark-recapture or sacrifice
510 methods): sex-determination system is significantly related to ASR in amphibians, reptiles
511 and tetrapods (PGLS results, amphibians: $b \pm SE = 0.09 \pm 0.03$, $t = 3.07$, $P = 0.004$, $n = 35$
512 species; reptiles: $b \pm SE = 0.11 \pm 0.03$, $t = 3.974$, $P < 0.001$, $n = 22$; tetrapods with snakes,
513 birds and mammals included as single data points: $b \pm SE = 0.10 \pm 0.02$, $t = 4.23$, $P < 0.001$, n
514 $= 55$).

Population genetic models

We developed population genetic models of the effects that deleterious mutation and sex-antagonistic selection might have on the ASR (Supplementary Material 2). The models assume that deleterious mutations are largely or entirely recessive, that they have multiplicative fitness effects across loci, that the loci are fully sex-linked and in linkage equilibrium, that mutation is not sex-biased, and that selection is strong relative to mutation and drift. Fitness effects of mutations in hemizygotes and homozygotes are assumed equal. Full details of the models are given in Supplementary Material 2. Here we summarize key results.

When deleterious alleles reach a mutation-selection balance, with X-Y sex determination the mean viability of males relative to females is

$$\overline{W}_m \approx \exp\{-3U_X - U_Y\},$$

where U_X and U_Y are the total rates of mutation to deleterious alleles across all loci on the X and Y chromosomes. With Z-W sex determination, the mean viability of females to males is

$$\overline{W}_f \approx \exp\{-3U_Z - U_W\},$$

where U_Z and U_W are the total rates of mutation to deleterious alleles across all loci on the Z and W chromosomes. Using very rough estimates for rates of deleterious mutations appropriate for human sex chromosomes, we estimate that mutation-selection balance might bias the ASR by a few percent. This degree of bias is substantially less than what is

seen in our data. We emphasize that the conclusion could be quite different using other parameter values, or if the model was extended to include stochastic effects.

The second hypothesis to explain biased ASRs that we explored with models is sex-antagonistic selection, the situation in which alleles are selected differently in females and males⁶⁷. In Supplementary Material 2, we use numerical examples to show that under both XY and ZW sex determination, either a female-biased or male-biased ASR can result. Thus there does not seem to be a robust generalization about how sex-antagonistic selection will bias the ASR.

Methods References

- 56 Jongepier, E. Reptilian adult sex ratios are biased towards the homogametic sex. Masters thesis, University of Groningen, 40 pp. (2011).
57. Arendt, J. D., Reznick, D. N. & López-Sepulcre, A. Replicated origin of female-biased adult sex ratio in introduced populations of the Trinidadian Guppy (*Poecilia reticulata*). *Evolution*, **68**, 2343–2356 (2014).
58. Freckleton, R. P., Harvey, P. H. & Pagel, M. Phylogenetic analysis and comparative data: a test and review of evidence. *Am. Nat.* **160**, 712–726 (2002).
59. Encyclopedia of Life. Available at <http://www.eol.org>. (2011). doi:10.4016/30633.0161.
60. Lislevand, T., Figuerola, J. & Székely, T. Avian body sizes in relation to fecundity, mating system, display behaviour and resource sharing. *Ecology* **88**, 1605 (2007).
61. Jones, K. E. *et al.* PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology* **90**, (2009).
62. Healy K. *et al.* Ecology and mode of life explain lifespan variation in birds and mammals. *Proc. R. Soc. B* **281** (2014).
63. Lessells, C. M. & Boag, P. T. Unrepeatable repeatabilities? A common mistake. *Auk* **104**, 116–121 (1987).
64. Maddison, W. P. & Maddison, D. R. Mesquite: a modular system for evolutionary analysis. (2011).

- 569 65. Pagel, M. & Meade, A. Bayesian analysis of correlated evolution of discrete characters
570 by reversible-jump Markov chain Monte Carlo. *Am. Nat.*, **167**, 808-825 (2006).
- 571 66. R Development Core Team. R: A language and environment for statistical computing.
572 R Foundation for Statistical Computing. Vienna, Austria. ISBN 3-900051-07-0, URL
573 <http://www.R-project.org> (2008).
- 574 67. Bennett, J. H. Selectively balanced polymorphism at a sex-linked locus. *Nature*, **180**,
575 1363–1364 (1957).

Extended Data titles and legends

Extended Data Table 1. Detailed analyses of the effect of sex-determination system on the adult sex ratio.

These are extensions of Table 1 in the text showing details of the phylogenetically uncorrected (t-tests) and phylogenetically corrected analyses (PGLS²⁴ and PDM²⁵). Birds and mammals were not tested with phylogenetic control because there is no variation in the type of sex-determination system within birds and mammals.

Extended Data Table2. Phylogenetically controlled analyses of the relationship between adult sex ratio and genetic sex-determination system using different branch length assumptions.

These are results of phylogenetic generalized least squares models (PGLS)²⁴ as implemented in the R package ‘caper’⁴⁶. The models assume gradual branch lengths calculated either by Nee’s or by Pagel’s method, or unit branch lengths⁶¹.

Extended Data Figure 1. Phylogenetically corrected mean and standard error of adult sex ratio in clades with different sex-determination systems.

Parameter estimates for means and the associated standard errors were calculated by phylogenetic generalized least squares models (PGLS)²⁴ presented in Extended Data Table 2 (with branch lengths estimated by Nee's method).